

## Article

# Effects of Meteorological Factors and Anthropogenic Precursors on PM<sub>2.5</sub> Concentrations in Cities in China

Ziyun Jing <sup>1,2</sup>, Pengfei Liu <sup>1,2,\*†</sup> , Tuanhui Wang <sup>2,3</sup>, Hongquan Song <sup>2,3,4,\*</sup>, Jay Lee <sup>5,6</sup>, Tao Xu <sup>7,8</sup> and Yu Xing <sup>9</sup>

- <sup>1</sup> Key Research Institute of Yellow River Civilization and Sustainable Development & Collaborative Innovation Center on Yellow River Civilization of Henan Province, Henan University, Kaifeng 475004, Henan, China; jzy@henu.edu.cn
  - <sup>2</sup> Institute of Urban Big Data, College of Environment and Planning, Henan University, Kaifeng 475004, Henan, China; thwang@henu.edu.cn
  - <sup>3</sup> Laboratory of Geospatial Technology for the Middle and Lower Yellow River Regions, Ministry of Education, Henan University, Kaifeng 475004, Henan, China
  - <sup>4</sup> Henan Key Laboratory of Integrated Air Pollution Control and Ecological Security, Henan University, Kaifeng 475004, Henan, China
  - <sup>5</sup> College of Environment and Planning, Henan University, Kaifeng 475004, Henan, China; jlee@kent.edu
  - <sup>6</sup> Department of Geography, Kent State University, Kent, OH 44242, USA
  - <sup>7</sup> School of Computer and Information Engineering, Henan University, Kaifeng 475004, Henan, China; txu@henu.edu.cn
  - <sup>8</sup> Henan Key Laboratory of Big Data Analysis and Processing, Henan University, Kaifeng 475004, Henan, China
  - <sup>9</sup> Henan Ecological and Environmental Monitoring Center, Zhengzhou 450000, Henan, China; nikkoyu@126.com
- \* Correspondence: lpf@henu.edu.cn (P.L.); hqsong@henu.edu.cn (H.S.)  
 † These authors contributed equally to this work and should be considered co-first authors.

Received: 7 February 2020; Accepted: 20 April 2020; Published: 27 April 2020



**Abstract:** Fine particulate matter smaller than 2.5  $\mu\text{m}$  (PM<sub>2.5</sub>) in size can significantly affect human health, atmospheric visibility, climate, and ecosystems. PM<sub>2.5</sub> has become the major air pollutant in most cities of China. However, influencing factors and their interactive effects on PM<sub>2.5</sub> concentrations remain unclear. This study used a geographic detector method to quantify the effects of anthropogenic precursors (AP) and meteorological factors on PM<sub>2.5</sub> concentrations in cities of China. Results showed that impacts of meteorological conditions and AP on PM<sub>2.5</sub> have significant spatio-temporal disparities. Temperature was the main influencing factor throughout the whole year, which can explain 27% of PM<sub>2.5</sub> concentrations. Precipitation and temperature were primary impacting factors in southern and northern China, respectively, at the annual time scale. In winter, AP had stronger impacts on PM<sub>2.5</sub> in northern China than in other seasons. Ammonia had stronger impacts on PM<sub>2.5</sub> than other anthropogenic precursors in winter. The interaction between all factors enhanced the formation of PM<sub>2.5</sub> concentrations. The interaction between ammonia and temperature had strongest impacts at the national scale, explaining 46% ( $q = 0.46$ ) of PM<sub>2.5</sub> concentrations. The findings comprehensively elucidated the relative importance of driving factors in PM<sub>2.5</sub> formation, which can provide basic foundations for understanding the meteorological and anthropogenic influences on the concentration patterns of PM<sub>2.5</sub>.

**Keywords:** PM<sub>2.5</sub>; particulate matter; interactions; **GeoDetector**; air quality

## 1. Introduction

With rapid urbanization and economic development, China has experienced increased emissions of greenhouse gases and atmospheric pollutants [1]. Particulate matter (PM), especially fine PM with a diameter less than  $2.5\ \mu\text{m}$  ( $\text{PM}_{2.5}$ ), has significant impacts on human health [2–6], atmospheric visibility [7,8], climate [9,10], and ecosystems [11,12]. In 190 Chinese cities, the annual mean  $\text{PM}_{2.5}$  concentration ( $57 \pm 18\ \mu\text{g m}^{-3}$ ) severely exceeds the limited value ( $35\ \mu\text{g m}^{-3}$ ) recommended by the Chinese ambient air quality standards (CAAQS) (GB3095-2012) [13], which is at least 1.6 times the limited value.  $\text{PM}_{2.5}$  pollution is a severe threat to the economic and social sustainability of China.

$\text{PM}_{2.5}$  concentrations could be affected by numerous factors such as meteorological conditions and anthropogenic precursors (AP) [14–17]. Meteorological conditions play important roles in the process of air pollutant emission, transport, dilution, chemical conversion, and deposition [18–20]. AP directly or indirectly affect  $\text{PM}_{2.5}$  concentrations in the atmosphere through a series of photochemical reactions under specific meteorological conditions [21–23]. Previous studies have explored the effects of AP and meteorological conditions (MCs) on the formation of  $\text{PM}_{2.5}$  concentrations. They found that temperature affects  $\text{PM}_{2.5}$  concentrations through atmospheric disturbances and photochemical reactions [24]. For example, as temperature and sulfur dioxide levels ( $\text{SO}_2$ ) rise, this leads to increased sulfate concentrations [25,26]. In addition, temperature has indirect effects on secondary organic aerosols. Under high-temperature conditions, higher biomass volatile organic compound (VOC) emissions can increase secondary organic aerosol concentrations [27]. Wind speed can change the diffusion and transport of  $\text{PM}_{2.5}$  in the horizontal direction [28,29]. Wet deposition caused by precipitation has scavenging effects on  $\text{PM}_{2.5}$  in most areas [25,29]. Absolute humidity also affects  $\text{PM}_{2.5}$  concentrations, and the increase in humidity is conducive to the conversion of semi-volatile substances into an aerosol phase, thereby increasing  $\text{PM}_{2.5}$  concentrations [30,31]. Moreover, the height of the mixing layer also plays an important role in the dilution of primary and secondary aerosols [32,33].

Due to the limitation of data accessibility before 2013 in China, research on  $\text{PM}_{2.5}$  was concentrated in Beijing-Tianjin-Hebei [17,34,35], the Yangtze River Delta [36–38], and the Pearl River Delta [39–42]. With the release of national air quality monitoring data, it is possible to study the temporal and spatial variations [43–45] and driving mechanisms of  $\text{PM}_{2.5}$  concentrations [46–48] throughout China. In recent years, some studies have been conducted to identify spatial distributions and temporal trends in  $\text{PM}_{2.5}$  concentrations at the national level, and to explore their relationships with meteorological conditions and precursor emissions. A series of models have been used to identify driving mechanisms influencing factors in  $\text{PM}_{2.5}$  formation, such as the classical ordinary least square model, geographically weighted regression model [49], spatial regression model [50], and land-use regression and other models [51]. The mechanisms of  $\text{PM}_{2.5}$  formation are very complicated, and there are interactions between influencing factors. The complex coupling of factors remains unclear. Under different driving conditions,  $\text{PM}_{2.5}$  concentrations have greater heterogeneity at different spatial and temporal scales. Moreover, traditional methods can lead to biased results because the impacting factors can interact closely with each other. Therefore, advanced methods should be adopted to explore the relative importance of the effects of each influencing factor on  $\text{PM}_{2.5}$ .

To solve this issue, we applied a geographical detector  $q$  statistical method (GeoDetector) to assess the influence of AP and MCs on the concentrations of  $\text{PM}_{2.5}$  in major cities of China. This study can provide a better understanding of the contribution of a single impacting factor on  $\text{PM}_{2.5}$  concentration. The findings can also provide a foundation for the improvement of  $\text{PM}_{2.5}$  prediction models' performance.

## 2. Materials and Methods

### 2.1. Study Area

Given differences in economic development, topography, and climate patterns, mainland China can be typically divided into 10 regions (Supplementary Materials, Figure S1). These 10 regions are

NE (northeastern region), NC (northern coastal region), EC (eastern coastal region), SC (southeastern coastal region), MUYR (Yangtze River's middle and upper reaches), MUPR (Pearl River's middle and upper reaches), UYR (Yellow River's upper reaches), MYR (Yellow River's middle reaches), XJ (Xinjiang region), and QTP (Qinghai Tibetan Plateau region) [29]. Among them, MUPR, SC, EC, and MUYR are considered to be southern China, while NC, NE, MYR, and UYR are considered to belong to northern China.

## 2.2. Datasets

According to CAAQS, annual average  $PM_{2.5}$  concentrations are limited to  $15 \mu g m^{-3}$  (Grade I) and  $35 \mu g m^{-3}$  (Grade II). The daily average concentrations are  $35 \mu g m^{-3}$  (Grade I) and  $75 \mu g m^{-3}$  (Grade II) [39]. Grade I refers to the concentration limit required for scenic spots, nature reserves, and other areas requiring special conservation in China. Grade II refers to the concentration limits required for rural areas, residential areas, industrial areas, cultural areas, and mixed-use residential areas. The daily  $PM_{2.5}$  concentrations of 366 cities in China were obtained from the China Environmental Monitoring Center. Due to the availability of data, data for Hong Kong, Macau, and Taiwan were not included. Figures S3 and S4 show maps of  $PM_{2.5}$  concentrations [29].

We acquired meteorological data (839 sites) from the China Meteorological Data Network throughout the whole year of 2016. The daily meteorological data included surface air pressure (PS, hPa), air temperature (TE, °C), relative humidity (RH, %), wind velocity (WI,  $m s^{-1}$ ), sunshine duration (SS, h), and accumulated precipitation (PE, mm) (Figures S5–S11) [29]. The monthly anthropogenic emissions of VOCs ammonia ( $NH_3$ ), sulfur dioxide ( $SO_2$ ), and nitrogen oxide ( $NO_x$ ) were monitored in 2016. They were usually considered as AP of  $PM_{2.5}$  and were collected from MEIC (multi-resolution emission inventory for China, <http://www.meicmodel.org/>). The MEIC includes emission data of the four sub-sectors of transportation, power, industry, and residential [52], and has been widely used in the research of air pollution [53–56]. Figure S12 indicates that the highest anthropogenic precursor emissions are mainly distributed in EC, MYR, SC, NC, and MUYR.

## 2.3. GeoDetector

In this study, the  $q$  statistics of GeoDetector were used to quantitatively analyze the impacts of AP and MCs on  $PM_{2.5}$  in China. GeoDetector supports a series of statistical methods that can explore spatial difference and identify the driving factors. The main idea is based on the assumption that if an independent variable ( $X$ ) causes a dependent variable ( $Y$ ), then the spatial distribution of the independent variable and the dependent variable should be consistent [57–60]. GeoDetector can detect both qualitative data and numerical data. Compared with traditional linear statistical methods, this is a major advantage of GeoDetector. Another unique advantage of GeoDetector is the ability to detect the interaction between two factors acting on the dependent variable. The GeoDetector includes four detectors, which are factor detection, risk area detection, ecological detection, and interaction detection. In this study, factor detection and interaction detection were used.

Factor detector uses  $q$  statistic to detect the influence of  $X$  (e.g., MCs and AP) on  $Y$  (e.g., the  $PM_{2.5}$  concentrations). The expression is:

$$q = 1 - \frac{\sum_{h=1}^L N_h \sigma_h^2}{N \sigma^2} = 1 - \frac{SSW}{SST},$$

$$SSW = \sum_{h=1}^L N_h \sigma_h^2, SST = N \sigma^2$$


In the formula,  $h = 1, \dots, L$ , which classifies  $X$  or  $Y$ ;  $N$  and  $N_h$  are the numbers of the whole region and categories in classification  $h$ ;  $\sigma^2$  and  $\sigma_h^2$  are the  $Y$  value of the whole region and the variance of strata  $h$ , respectively.  $SST$  and  $SSW$  are the total variance of the whole region and the sum of variance within the strata, respectively. Greater values of  $q$  (0–1) indicate more spatial variation in  $Y$ . If the classification is based on  $X$ , a higher  $q$  value explains the influence of  $X$  on  $Y$  (i.e., explaining power:  $100 \times q\%$ ).

Interaction detection can identify the impact of the interaction between potential driving factors. Based on that, we can assess whether the interaction between  $X_1$  and  $X_2$  will strengthen or weaken the explaining power of  $Y$ . Additionally, the influences of these factors on the dependent variable  $Y$  would be independent of each other. There are five types of interactions; please refer to [59] for more information. In addition, in order to identify the positive or negative correlations between  $PM_{2.5}$  concentrations and influencing factors, this study calculated their Pearson correlation coefficients at different temporal and spatial scales.

### 3. Results

#### 3.1. Effects on $PM_{2.5}$ Concentrations at the National Scale

The influence of each driving factor on  $PM_{2.5}$  concentrations was acquired by calculating the corresponding  $q$  value (the power of determinant, Figure 1a), which indicated the contribution of each impacting factor on  $PM_{2.5}$  concentrations. Figure 1 shows that there were obvious seasonal and annual difference in factors' impacts on  $PM_{2.5}$ . Meteorological conditions were dominant impacting factors in  $PM_{2.5}$  formation at the annual time scale. TE ( $q = 0.27$ ) was the primary impacting factor, followed by PE ( $q = 0.22$ ) and PS ( $q = 0.17$ ).




**Figure 1.** The annual and seasonal  $q$  values (a) and correlation coefficient (b) between  $PM_{2.5}$  concentrations and influencing factors at the national scale. Note: VO denotes volatile organic compounds (VOCs); NO denotes  $NO_x$ ; NH denotes  $NH_3$ ; SO denotes  $SO_2$ .

Meteorological factors were dominant driving forces in spring, such as PE ( $q = 0.12$ ) and PS ( $q = 0.10$ ). In summer, AP showed stronger impacts on  $PM_{2.5}$  concentrations than meteorological conditions, and the  $NO_x$ ,  $NH_3$ , and VOCs were the three dominant factors ( $q > 0.10$ ). In autumn, the meteorological factors and AP showed comparative influence on  $PM_{2.5}$ . The dominant impacting factor in autumn was TE ( $q = 0.18$ ), followed by PE ( $q = 0.13$ ),  $NO_x$  ( $q = 0.11$ ), and  $NH_3$  ( $q = 0.10$ ). Similar to the autumn, meteorological factors and AP had comparative impacts on  $PM_{2.5}$  concentrations in winter. TE ( $q = 0.25$ ) was the dominant factor, followed by  $NH_3$  ( $q = 0.18$ ), SSD ( $q = 0.17$ ), PS ( $q = 0.16$ ), and VOCs ( $q = 0.13$ ). This indicated that AP were the dominant impacting factor in winter.

### 3.2. Effects on $PM_{2.5}$ Concentrations at the Regional Scale

Figure 2 shows the effects of AP and MCs on  $PM_{2.5}$  significantly varied at regional and seasonal scales in China. In general, meteorological factors were the major driving forces in China. PE and TE were primary driving forces in southern and northern China, respectively.



**Figure 2.** The  $q$  values of impacting factors at the annual and seasonal time scales in 10 regions of China.  
Note: VO denotes VOCs; NO denotes  $NO_x$ ; NH denotes  $NH_3$ ; SO denotes  $SO_2$ .


Figure 2 shows that meteorological factors were primary drivers of  $PM_{2.5}$  formation in spring, which is similar to most regions at the annual time scale. The dominant meteorological factor was PE and SS in southern and regions of MYR and NC, respectively. Meteorological factors and AP showed comparative impacts on  $PM_{2.5}$  concentrations in summer except for in XJ and QTP. TE was the dominant driving factor on  $PM_{2.5}$  concentrations in autumn in UYR, NE, and MYR, but PE and WI played the dominant role in regions of MUYR and SC, and of NC and EC, respectively. WI was the primary driving factor in NC, NE, EC, and UYR in winter, but PS was the major driving factor in MUYR and MUPR. However, in QTP, PS was the major impacting factor on  $PM_{2.5}$  concentrations throughout the whole year.

### 3.3. Interactive Effects on $PM_{2.5}$

This study explored interactive effects on  $PM_{2.5}$  by using the interaction detector with a total of 45 pairs of interactions. The interaction of any two factors was analyzed by comparing their combined contribution with their individual contributions to  $PM_{2.5}$  concentrations. Figure 3 shows the  $q$  values of each pair of impact factors and their interaction through the whole year at the national scale. Interactions of  $PE \cap TE$ ,  $PS \cap TE$ ,  $PS \cap VO$ ,  $PS \cap NO$ ,  $VO \cap NO$ ,  $VO \cap NH$ , and  $NO \cap NH$  belong to bivariate enhancements and other interactions belong to nonlinear enhancements (Figure 3). Generally, the interaction between NH and TE ( $q$  value = 0.46) was the strongest interaction among all impacting factors. Figure S14 indicates that there were obvious seasonal disparities in the interactive influence. In spring, fall, and winter (but not summer), the interactions between meteorological factors played




major roles in  $PM_{2.5}$  concentrations. The interaction  $WI \cap RH$  ( $q$  value = 0.38) had the strongest effect on  $PM_{2.5}$  in spring. In autumn and winter, the interaction between SS and TE (autumn:  $q$  value = 0.54; winter:  $q$  value = 0.42) played the strongest role in  $PM_{2.5}$  concentrations. However,  $NH \cap RH$  ( $q$  value = 0.35) was the highest in summer.



**Figure 3.** The annual interactive and individual  $q$  value of impacting factors in China. Note: VO denotes VOCs; NO denotes  $NO_x$ ; NH denotes  $NH_3$ ; SO denotes  $SO_2$ .

There were obvious regional disparities in all the interactions in China (Figure 4). The interactions between meteorological factors were strongest in all regions. In the interactions between AP and meteorological factors, the interaction of  $AP \cap TE$  played a primary role in EC, MUYR, and northern China, and the interaction of  $AP \cap PE$  played a dominant role in MUYR, MUPR, and SC. Figures S15–S18 show the seasonal interactive effects of each pair of driving factors at the regional scale in China, indicating that there were also obvious regional and seasonal differences in all the interactions.



**Figure 4.** Annual interactions between driving factors at the regional scale (we listed the top 15 interactions). Note: VO denotes VOCs; NO denotes  $NO_x$ ; NH denotes  $NH_3$ ; SO denotes  $SO_2$ .

#### 4. Discussion

PM<sub>2.5</sub> concentrations were found to be driven by natural conditions and human activities. Previous studies had reported that PM<sub>2.5</sub> concentrations correlate with both MCs and AP emissions [16,61–65]. We analyzed influences of AP and meteorological factors and their interactions on PM<sub>2.5</sub> in Chinese cities. Results showed that effects of AP and MCs and their interactions on PM<sub>2.5</sub> had obvious seasonal and regional variations across China.

In this study, we found that meteorological factors were the leading driving factors at the annual time scale and the national scale, indicating that the meteorological conditions had dominant influences on PM<sub>2.5</sub> concentrations. TE and PE were the two leading factors affecting PM<sub>2.5</sub> concentrations. TE was closely related to PM<sub>2.5</sub> concentrations by affecting atmospheric perturbation and chemical reactions [24]. PE could scavenge PM<sub>2.5</sub> from the air and had a moisture removal effect [11,66,67]. However, the major influencing factors on PM<sub>2.5</sub> concentration had significantly seasonal variations. The influences of meteorological conditions on PM<sub>2.5</sub> concentrations were the strongest in winter, and TE was the leading factor affecting PM<sub>2.5</sub> concentrations. This is because temperature inversion can weaken the scattering and dispersion of PM<sub>2.5</sub>, resulting in higher local PM<sub>2.5</sub> pollution [68,69]. In summer, influences of meteorological conditions on PM<sub>2.5</sub> concentrations were weakest. RH was the leading factor affecting PM<sub>2.5</sub> concentrations in summer. This was because RH was higher in summer than in the other three seasons, which had a suppression effect on PM<sub>2.5</sub> under moist air conditions [29].

At the annual time scale, TE was the dominant factor in northern and southern China (except MUPR), which is consistent with previous studies [70,71]. PE was the primary factor affecting PM<sub>2.5</sub> concentrations in southern China in most seasons throughout the whole year. This was because PE in southern China was higher than in other regions [29]. The increasing PE had scavenging effects on PM<sub>2.5</sub> by wet deposition, and could lower the PM<sub>2.5</sub> concentration [66,67]. WI was the main influencing factor in EC and northern China in winter, which is similar to some reports that WI was the most important and negative impacting factor on PM concentrations [72,73]. This was because weaker East Asia winter monsoons could slow wind speeds and increased the frequency of static wind, which had made it more difficult for PM to disperse [74].

Previous studies have indicated that AP were also crucial driving factors on PM<sub>2.5</sub> [48,63,75]. We found that the influence of AP on PM<sub>2.5</sub> in winter were higher than in other seasons in XJ and northern China (MYR, UYR, NC). This might be due to the high anthropogenic emissions from winter carbon-fired heating and less surface vegetation cover in winter in northern China, which significantly increased pollutant emissions in the atmosphere [76,77]. In addition, NH<sub>3</sub> had important impacts on PM<sub>2.5</sub> among AP factors in winter at the national scale and in some individual regions (NC, SC, MUJR, and MYR). Ammonia is an important precursor, and emissions of ammonia had stronger associations with PM<sub>2.5</sub> concentrations than other anthropogenic precursors; this is similar to results of previous studies [78]. Ammonia participates in photochemical reactions as an atmospheric alkaline gas, which is important in the SIA (secondary inorganic aerosol) formation of compounds such as ammonium salts, sulfate, and nitrate [79–83]. In China, SIA was an important driving factor of PM<sub>2.5</sub> pollution, especially during severe smog events [80,84]. SIA accounted for 32% of PM<sub>2.5</sub> mass concentration in China (e.g., Beijing, Guangzhou, Shanghai, and Xi'an) during the 2013 haze pollution events [84]. There was a 5.7% reduction in the annual concentration of PM<sub>2.5</sub> concentrations when ammonia emissions were cut by 47% in the Beijing-Tianjin-Hebei region of China [79].

PM<sub>2.5</sub> concentrations are affected by complex interactions between AP and meteorological conditions. This study found that interactions between any driving factors at all time and space scales had significant enhancement effects on PM<sub>2.5</sub> concentrations. The leading interactive effect between AP and MCs was between AP with RH and PE at the national scale in summer. This might be due to the fact that PE and RH have scavenging and suppression effects on PM<sub>2.5</sub> by wet deposition in summertime. However, the primary interactive effect between AP and MCs was AP  $\cap$  TE at the national scale in winter. This was due to increased temperature inversion under the lower winter temperature, which weakened the diffusion and dispersion of pollution. The interaction between

AP had the dominant effect on  $PM_{2.5}$  concentrations over most regions in summer, which indicated that accelerated photochemical reactions between AP occurred under high-temperature conditions. Interactions between AP and factors of PE and PS were important at annual and seasonal scales in southern China. This might be due to the high precipitation and surface pressure in southern China [29].

## 5. Conclusions

The effects of AP and MCs on  $PM_{2.5}$  in Chinese cities were systematically analyzed in this study. The findings revealed significant seasonal and regional disparities in the impacts of examined factors and how they interacted on  $PM_{2.5}$ . The results can help us to better understanding the relative importance of the driving factors in the formation of  $PM_{2.5}$ . The study indicated that local AP and meteorological factors had important impacts on  $PM_{2.5}$  in China, and had obvious regional and seasonal variations. Meteorological conditions played a leading role in determining  $PM_{2.5}$  concentrations at the regional and national scales throughout the whole year. At the seasonal time scale, WI was the primary factor on  $PM_{2.5}$  concentrations in winter in northern China and XJ, but PE was the major driver on  $PM_{2.5}$  concentrations during most seasons in southern China. However, AP had stronger impacts on  $PM_{2.5}$  during winter than in other seasons in XJ and most regions of northern China.  $NH_3$  had a stronger effect on  $PM_{2.5}$  concentrations during winter than other anthropogenic precursors. Interactions between all influencing factors have enhanced effects on  $PM_{2.5}$  concentrations. In addition, the interaction between MCs and AP played a leading role at the national scale throughout the whole year and in summer and winter. The results could provide a basis for the government to develop more precise air pollution control strategies.

Some limitations to the study should be clarified to assist future studies. First, land use and land cover, socioeconomic conditions, elevation, and topography were not considered to assess their influence on  $PM_{2.5}$  concentrations. Second, the uncertainty of MEIC emission inventory may lead to some uncertainties in the research results. When processing emissions inventory data, because the inventory only has monthly emissions data, so as to match the daily  $PM_{2.5}$  concentration and daily weather monitoring data, we took the arithmetic average of the monthly inventory emissions data, which also increased the uncertainty of the analysis results. Third, the data used in this study is limited to 2016, and does not include data analysis of other years. This is because the emission inventory data that we could obtain were for 2008, 2010, 2012, 2014, and 2016, but the  $PM_{2.5}$  data in 366 cities were available only from 2015 to 2017. In order to maintain consistency between the data, we selected 2016 as the research period in this study. There was no comparative analysis of inter-annual variability. Therefore, we should comprehensively consider other factors on  $PM_{2.5}$  concentrations including socioeconomic, land use, terrain, and elevation in the future. In addition, inter-annual change analysis based on multi-year data needs to be added.

**Supplementary Materials:** The following are available online at <http://www.mdpi.com/2071-1050/12/9/3550/s1>, Figure S1: Ten regions in China, including NE, NC, EC, SC, MYR, MUYR, MUPR, UYR, XJ, and QTP. Figure S2: Map of air quality monitoring sites (a) and meteorological sites (b). Figure S3: Annual mean  $PM_{2.5}$  concentrations in Chinese cities (2015–2017). Figure S4: Seasonal mean  $PM_{2.5}$  concentrations in spring (a), summer (b), autumn (c), and winter (d) in Chinese cities (2015–2017). Figure S5: Spatial distributions of annual mean precipitation (a), surface pressure (b), relative humidity (c), sunshine duration (d), air temperature (e), and wind speed (f) in China during 2015–2017. Figure S6: Spatial distributions of seasonal precipitations in spring (a), summer (b), autumn (c), and winter (d) in China during 2015–2017. Figure S7: Spatial distributions of seasonal surface pressure in spring (a), summer (b), autumn (c), and winter (d) in China during 2015–2017. Figure S8: Spatial distributions of seasonal relative humidity in spring (a), summer (b), autumn (c), and winter (d) in China during 2015–2017. Figure S9: Spatial distributions of seasonal sunshine duration in spring (a), summer (b), autumn (c), and winter (d) in China during 2015–2017. Figure S10: Spatial distributions of seasonal air temperature in spring (a), summer (b), autumn (c), and winter (d) in China during 2015–2017. Figure S11: Spatial distributions of seasonal wind speed in spring (a), summer (b), autumn (c), and winter (d) in China during 2015–2017. Figure S12: (a)  $NH_3$  emissions, (b)  $NO_x$  emissions, (c)  $SO_2$  emissions, and (d) VOCs in China in 2016 with resolution of  $0.25^\circ \times 0.25^\circ$ . Figure S13: Correlations between  $PM_{2.5}$  concentrations and impacting factors in 10 regions (a. EC, b. MUPR, c. MUYR, d. MYR, e. NC, f. NE, g. QTP, h. SC, i. UYR, and j. XJ) of China. Figure S14: The seasonal interactive  $q$  values and the original  $q$  value of each pair of factors. Figure S15: The interactions between impacting factors in spring at the



regional scale in China. Figure S16: Interactions between impacting factors in summer at the regional scale in China. Figure S17: Interactions between impacting factors in autumn at the regional scale in China. Figure S18: Interactions between impacting factors in winter at the regional scale in China. Table S1: Effect of various factors on PM<sub>2.5</sub> in China in 2016. Table S2: Effect of various factors on PM<sub>2.5</sub> throughout the whole year at the regional scale. Table S3: Effect of various factors on PM<sub>2.5</sub> in spring at the regional scale. Table S4: Effect of various factors on PM<sub>2.5</sub> in summer at the regional scale. Table S5: Effect of various factors on PM<sub>2.5</sub> in autumn at the regional scale. Table S6: Effect of various factors on PM<sub>2.5</sub> in winter at the regional scale.

**Author Contributions:** Conceptualization, H.S.; methodology, H.S.; software, P.L., T.X.; validation, P.L., T.W. and J.L.; formal analysis, P.L.; investigation, P.L.; resources, H.S. and Y.X.; data curation, P.L.; writing—original draft preparation, Z.J., P.L.; writing—review and editing, P.L. and H.S.; visualization, T.W.; supervision, H.S. and J.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Natural Science Foundation of China (41401107) and the Basic and Frontier Technology Research Project of Henan Province, China (162300410132), and the Higher Education Research Project of Henan Province, China (17B170003).

**Acknowledgments:** We the authors thank Shenghui Zhou from Henan University for helpful discussions on data analysis.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Liu, X.H.; Zhang, Y.; Xing, J.; Zhang, Q.; Wang, K.; Streets, D.G.; Jang, C.; Wang, W.X.; Hao, J.M. Understanding of regional air pollution over China using CMAQ, part II. Process analysis and sensitivity of ozone and particulate matter to precursor emissions. *Atmos. Environ.* **2010**, *44*, 3719–3727. [[CrossRef](#)]
2. Zhang, Q.; He, K.; Hong, H. Cleaning China's air. *Nature* **2012**, *484*, 161–162. [[CrossRef](#)] [[PubMed](#)]
3. Watson, T. Environment: Breathing trouble. *Nature* **2014**, *513*, S14–S15. [[CrossRef](#)] [[PubMed](#)]
4. Jerrett, M. Atmospheric science: The death toll from air-pollution sources. *Nature* **2015**, *525*, 330–331. [[CrossRef](#)] [[PubMed](#)]
5. Kerl, P.Y.; Zhang, W.; Moreno-Cruz, J.B.; Nenes, A.; Realff, M.J.; Russell, A.G.; Sokol, J.; Thomas, V.M. New approach for optimal electricity planning and dispatching with hourly time-scale air quality and health considerations. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 10884–10889. [[CrossRef](#)] [[PubMed](#)]
6. Patz, J.A.; Campbell-Lendrum, D.; Holloway, T.; Foley, J.A. Impact of regional climate change on human health. *Nature* **2005**, *438*, 310–317. [[CrossRef](#)] [[PubMed](#)]
7. Pui, D.Y.H.; Chen, S.-C.; Zuo, Z. PM<sub>2.5</sub> in China: Measurements, sources, visibility and health effects, and mitigation. *Particuology* **2014**, *13*, 1–26. [[CrossRef](#)]
8. Liu, Y.-J.; Zhang, T.-T.; Liu, Q.-Y.; Zhang, R.-J.; Sun, Z.-Q.; Zhang, M.-G. Seasonal variation of physical and chemical properties in TSP, PM<sub>10</sub> and PM<sub>2.5</sub> at a roadside site in Beijing and their influence on atmospheric visibility. *Aerosol Air Qual. Res.* **2014**, *14*, 954–969. [[CrossRef](#)]
9. Song, H.; Zhang, K.; Piao, S.; Wan, S. Spatial and temporal variations of spring dust emissions in northern China over the last 30 years. *Atmos. Environ.* **2016**, *126*, 117–127. [[CrossRef](#)]
10. Song, H.; Wang, K.; Zhang, Y.; Hong, C.; Zhou, S. Simulation and evaluation of dust emissions with WRF-Chem (v3.7.1) and its relationship to the changing climate over East Asia from 1980 to 2015. *Atmos. Environ.* **2017**, *167*, 511–522. [[CrossRef](#)]
11. Song, H.; Zhang, K.; Piao, S.; Liu, L.; Wang, Y.-P.; Chen, Y.; Yang, Z.; Zhu, L.; Wan, S. Soil organic carbon and nutrient losses resulted from spring dust emissions in Northern China. *Atmos. Environ.* **2019**, *213*, 585–596. [[CrossRef](#)]
12. Liu, J.; Rühland, K.M.; Chen, J.; Xu, Y.; Chen, S.; Chen, Q.; Huang, W.; Xu, Q.; Chen, F.; Smol, J.P. Aerosol-weakened summer monsoons decrease lake fertilization on the Chinese Loess Plateau. *Nat. Clim. Chang.* **2017**, *7*, 190–194. [[CrossRef](#)]
13. Zhang, Y.-L.; Cao, F. Fine particulate matter (PM<sub>2.5</sub>) in China at a city level. *Sci. Rep.* **2015**, *5*, 14884–14896. [[CrossRef](#)] [[PubMed](#)]
14. Tai, A.P.K.; Mickley, L.J.; Jacob, D.J. Correlations between fine particulate matter (PM<sub>2.5</sub>) and meteorological variables in the United States: Implications for the sensitivity of PM<sub>2.5</sub> to climate change. *Atmos. Environ.* **2010**, *44*, 3976–3984. [[CrossRef](#)]

15. Yang, D.; Wang, X.; Xu, J.; Xu, C.; Lu, D.; Ye, C.; Wang, Z.; Bai, L. Quantifying the influence of natural and socioeconomic factors and their interactive impact on PM<sub>2.5</sub> pollution in China. *Environ. Pollut.* **2018**, *241*, 475–483. [\[CrossRef\]](#)
16. Zhou, C.; Chen, J.; Wang, S. Examining the effects of socioeconomic development on fine particulate matter (PM<sub>2.5</sub>) in China's cities using spatial regression and the geographical detector technique. *Sci. Total Environ.* **2018**, *619–620*, 436–445. [\[CrossRef\]](#)
17. Gui, K.; Che, H.; Wang, Y.; Wang, H.; Zhang, L.; Zhao, H.; Zheng, Y.; Sun, T.; Zhang, X. Satellite-derived PM<sub>2.5</sub> concentration trends over Eastern China from 1998 to 2016: Relationships to emissions and meteorological parameters. *Environ. Pollut.* **2019**, *247*, 1125–1133. [\[CrossRef\]](#)
18. Hogrefe, C. Air quality: Emissions versus climate change. *Nat. Geosci.* **2012**, *5*, 685–686. [\[CrossRef\]](#)
19. Kinney, P.L. Climate Change, Air Quality, and Human Health. *Am. J. Prev. Med.* **2008**, *35*, 459–467. [\[CrossRef\]](#)
20. Lacressonnière, G.; Foret, G.; Beekmann, M.; Siour, G.; Engardt, M.; Gauss, M.; Watson, L.; Andersson, C.; Colette, A.; Josse, B.; et al. Impacts of regional climate change on air quality projections and associated uncertainties. *Clim. Chang.* **2016**, *136*, 309–324. [\[CrossRef\]](#)
21. Chang, W.; Zhan, J.; Zhang, Y.; Li, Z.; Xing, J.; Li, J. Emission-driven changes in anthropogenic aerosol concentrations in China during 1970–2010 and its implications for PM<sub>2.5</sub> control policy. *Atmos. Res.* **2018**, *212*, 106–119. [\[CrossRef\]](#)
22. Arif, M.; Kumar, R.; Kumar, R.; Eric, Z.; Gourav, P. Ambient black carbon, PM<sub>2.5</sub> and PM<sub>10</sub> at Patna: Influence of anthropogenic emissions and brick kilns. *Sci. Total Environ.* **2018**, *624*, 1387–1400. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Blanchard, C.L.; Shaw, S.L.; Edgerton, E.S.; Schwab, J.J. Emission influences on air pollutant concentrations in New York state: II. PM<sub>2.5</sub> organic and elemental carbon constituents. *Atmos. Environ. X* **2019**, *3*, 100039. [\[CrossRef\]](#)
24. He, J.; Gong, S.; Yu, Y.; Yu, L.; Wu, L.; Mao, H.; Song, C.; Zhao, S.; Liu, H.; Li, X.; et al. Air pollution characteristics and their relation to meteorological conditions during 2014–2015 in major Chinese cities. *Environ. Pollut.* **2017**, *223*, 484–496. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Lecoœur, E.; Seigneur, C. Dynamic evaluation of a multi-year model simulation of particulate matter concentrations over Europe. *Atmos. Chem. Phys.* **2013**, *13*, 4319–4337. [\[CrossRef\]](#)
26. Jacob, D.J.; Winner, D.A. Effect of climate change on air quality. *Atmos. Environ.* **2009**, *43*, 51–63. [\[CrossRef\]](#)
27. Heald, C.L.; Henze, D.K.; Horowitz, L.W.; Feddema, J.; Lamarque, J.F.; Guenther, A.; Hess, P.G.; Vitt, F.; Seinfeld, J.H.; Godstein, A.H.; et al. Predicted change in global secondary organic aerosol concentrations in response to future climate, emissions, and land use change. *J. Geophys. Res. Atmos.* **2008**, *113*, 1–16. [\[CrossRef\]](#)
28. Aksoyoglu, S.; Keller, J.; Barmadimos, I.; Oderbolz, D.; Lanz, V.A.; Prévôt, A.S.H.; Baltensperger, U. Aerosol modelling in Europe with a focus on Switzerland during summer and winter episodes. *Atmos. Chem. Phys.* **2011**, *11*, 7355–7373. [\[CrossRef\]](#)
29. Li, X.; Song, H.; Zhai, S.; Lu, S.; Kong, Y.; Xia, H.; Zhao, H. Particulate matter pollution in Chinese cities: Areal-temporal variations and their relationships with meteorological conditions (2015–2017). *Environ. Pollut.* **2019**, *246*, 11–18. [\[CrossRef\]](#)
30. Dawson, J.P.; Racherla, P.N.; Lynn, B.H.; Adams, P.J.; Pandis, S.N. Impacts of climate change on regional and urban air quality in the eastern United States: Role of meteorology. *J. Geophys. Res.* **2009**, *114*, 1–11. [\[CrossRef\]](#)
31. Galindo, N.; Varea, M.; Gil-Moltó, J.; Yubero, E.; Nicolás, J. The Influence of Meteorology on Particulate Matter Concentrations at an Urban Mediterranean Location. *Water Air Soil Pollut.* **2011**, *215*, 365–372. [\[CrossRef\]](#)
32. Sandeep, A.; Rao, T.N.; Ramkiran, C.N.; Rao, S.V.B. Differences in Atmospheric Boundary-Layer Characteristics Between Wet and Dry Episodes of the Indian Summer Monsoon. *Bound. -Layer Meteorol.* **2014**, *153*, 217–236. [\[CrossRef\]](#)
33. Pay, M.T.; Jiménez-Guerrero, P.; Baldasano, J.M. Assessing sensitivity regimes of secondary inorganic aerosol formation in Europe with the CALIOPE-EU modeling system. *Atmos. Environ.* **2012**, *51*, 146–164. [\[CrossRef\]](#)
34. Yan, S.; Cao, H.; Chen, Y.; Wu, C.; Hong, T.; Fan, H. Spatial and temporal characteristics of air quality and air pollutants in 2013 in Beijing. *Environ. Sci. Pollut. Res.* **2016**, *23*, 13996–14007. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Chen, Z.; Xu, B.; Cai, J.; Gao, B. Understanding temporal patterns and characteristics of air quality in Beijing: A local and regional perspective. *Atmos. Environ.* **2016**, *127*, 303–315. [\[CrossRef\]](#)

36. Yang, D.; Lu, D.; Xu, J.; Ye, C.; Zhao, J.; Tian, G.; Wang, X.; Zhu, N. Predicting spatio-temporal concentrations of PM<sub>2.5</sub> using land use and meteorological data in Yangtze River Delta, China. *Stoch. Environ. Res. Risk Assess.* **2017**, *32*, 2445–2456. [\[CrossRef\]](#)
37. Lu, D.; Xu, J.; Yang, D.; Zhao, J. Spatio-temporal variation and influence factors of PM<sub>2.5</sub> concentrations in China from 1998 to 2014. *Atmos. Pollut. Res.* **2017**, *8*, 1151–1159. [\[CrossRef\]](#)
38. Ma, T.; Duan, F.; He, K.; Qin, Y.; Tong, D.; Geng, G.; Liu, X.; Li, H.; Yang, S.; Ye, S.; et al. Air pollution characteristics and their relationship with emissions and meteorology in the Yangtze River Delta region during 2014–2016. *J. Environ. Sci. (China)* **2019**, *83*, 8–20. [\[CrossRef\]](#)
39. Lai, S.; Zhao, Y.; Ding, A.; Zhang, Y.; Song, T.; Zheng, J.; Ho, K.F.; Lee, S.; Zhong, L. Characterization of PM<sub>2.5</sub> and the major chemical components during a 1-year campaign in rural Guangzhou, Southern China. *Atmos. Res.* **2016**, *167*, 208–215. [\[CrossRef\]](#)
40. Wang, Q.Q.; Huang, X.H.H.; Zhang, T.; Zhang, Q.; Feng, Y.; Yuan, Z.; Wu, D.; Lau, A.K.H.; Yu, J.Z. Organic tracer-based source analysis of PM<sub>2.5</sub> organic and elemental carbon: A case study at Dongguan in the Pearl River Delta, China. *Atmos. Environ.* **2015**, *118*, 164–175. [\[CrossRef\]](#)
41. Zhao, X.; Wang, X.; Ding, X.; He, Q.; Zhang, Z.; Liu, T.; Fu, X.; Gao, B.; Wang, Y.; Zhang, Y.; et al. Compositions and sources of organic acids in fine particles (PM<sub>2.5</sub>) over the Pearl River Delta region, south China. *J. Environ. Sci.* **2014**, *26*, 110–121. [\[CrossRef\]](#)
42. Huang, H.; Ho, K.F.; Lee, S.C.; Tsang, P.K.; Ho, S.S.H.; Zou, C.W.; Zou, S.C.; Cao, J.J.; Xu, H.M. Characteristics of carbonaceous aerosol in PM<sub>2.5</sub>: Pearl Delta River Region, China. *Atmos. Res.* **2012**, *104–105*, 227–236. [\[CrossRef\]](#)
43. Li, Y.; Dai, Z.; Liu, X. Analysis of Spatial-Temporal Characteristics of the PM<sub>2.5</sub> Concentrations in Weifang City, China. *Sustainability* **2018**, *10*, 2960. [\[CrossRef\]](#)
44. Jin, Q.; Fang, X.; Wen, B.; Shan, A. Spatio-temporal variations of PM<sub>2.5</sub> emission in China from 2005 to 2014. *Chemosphere* **2017**, *183*, 429–436. [\[CrossRef\]](#) [\[PubMed\]](#)
45. Fang, C.; Wang, Z.; Xu, G. Spatial-temporal characteristics of PM<sub>2.5</sub> in China: A city-level perspective analysis. *J. Geogr. Sci.* **2016**, *26*, 1519–1532. [\[CrossRef\]](#)
46. Wang, S.; Liu, X.; Yang, X.; Zou, B.; Wang, J. Spatial variations of PM<sub>2.5</sub> in Chinese cities for the joint impacts of human activities and natural conditions: A global and local regression perspective. *J. Clean. Prod.* **2018**, *203*, 143–152. [\[CrossRef\]](#)
47. Wang, S.; Zhou, C.; Wang, Z.; Feng, K.; Hubacek, K. The characteristics and drivers of fine particulate matter (PM<sub>2.5</sub>) distribution in China. *J. Clean. Prod.* **2017**, *142*, 1800–1809. [\[CrossRef\]](#)
48. Wang, P.; Guo, H.; Hu, J.; Kota, S.H.; Ying, Q.; Zhang, H. Responses of PM<sub>2.5</sub> and O<sub>3</sub> concentrations to changes of meteorology and emissions in China. *Sci. Total Environ.* **2019**, *662*, 297–306. [\[CrossRef\]](#)
49. Hu, X.; Waller, L.A.; Al-Hamdan, M.Z.; Crosson, W.L.; Estes, M.G.; Estes, S.M.; Quattrochi, D.A.; Sarnat, J.A.; Liu, Y. Estimating ground-level PM<sub>2.5</sub> concentrations in the southeastern U.S. using geographically weighted regression. *Environ. Res.* **2013**, *121*, 1–10. [\[CrossRef\]](#)
50. Hao, Y.; Liu, Y.M. The influential factors of urban PM<sub>2.5</sub> concentrations in China: A spatial econometric analysis. *J. Clean. Prod.* **2016**, *112*, 1443–1453.
51. Mao, L.; Qiu, Y.; Kusano, C.; Xu, X. Predicting regional space–time variation of PM<sub>2.5</sub> with land-use regression model and MODIS data. *Environ. Sci. Pollut. Res.* **2012**, *19*, 128–138. [\[CrossRef\]](#)
52. Zhang, Q.; Streets, D.G.; Carmichael, G.R.; He, K.B.; Huo, H.; Kannari, A.; Klimont, Z.; Park, I.S.; Reddy, S.; Fu, J.S.; et al. Asian emissions in 2006 for the NASA INTEX-B mission. *Atmos. Chem. Phys.* **2009**, *9*, 5131–5153. [\[CrossRef\]](#)
53. Geng, G.; Zhang, Q.; Tong, D.; Li, M.; Zheng, Y.; Wang, S.; He, K. Chemical composition of ambient PM<sub>2.5</sub> over China and relationship to precursor emissions during 2005–2012. *Atmos. Chem. Phys.* **2017**, *17*, 9187–9203. [\[CrossRef\]](#)
54. Zheng, Y.; Zhang, Q.; Liu, Y.; Geng, G.; He, K. Estimating ground-level PM<sub>2.5</sub> concentrations over three megalopolises in China using satellite-derived aerosol optical depth measurements. *Atmos. Environ.* **2016**, *124*, 232–242. [\[CrossRef\]](#)
55. Jiang, X.; Hong, C.; Zheng, Y.; Zheng, B.; Guan, D.; Gouldson, A.; Zhang, Q.; He, K. To what extent can China’s near-term air pollution control policy protect air quality and human health? A case study of the Pearl River Delta region. *Environ. Res. Lett.* **2015**, *10*, 1–14. [\[CrossRef\]](#)

56. Zheng, B.; Zhang, Q.; Zhang, Y.; He, K.B.; Wang, K.; Zheng, G.J.; Duan, F.K.; Ma, Y.L.; Kimoto, T. Heterogeneous chemistry: A mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China. *Atmos. Chem. Phys. Discuss.* **2014**, *14*, 16731–16776. [[CrossRef](#)]
57. Wang, J.; Xu, C. Geodetector: Principle and prospective. *Acta Geogr. Sin.* **2017**, *72*, 116–134.
58. Wang, J.; Li, X.; Christakos, G.; Liao, Y.; Zhang, T.; Gu, X.; Zheng, X. Geographical Detectors-Based Health Risk Assessment and its Application in the Neural Tube Defects Study of the Heshun Region, China. *Int. J. Geogr. Inf. Sci.* **2010**, *24*, 107–127. [[CrossRef](#)]
59. Wang, J.F.; Zhang, T.L.; Fu, B.J. A measure of spatial stratified heterogeneity. *Ecol. Indic.* **2016**, *67*, 250–256. [[CrossRef](#)]
60. Wang, J.-F.; Hu, Y. Environmental health risk detection with GeogDetector. *Environ. Model. Softw.* **2012**, *33*, 114–115. [[CrossRef](#)]
61. Luo, J.; Du, P.; Samat, A.; Xia, J.; Che, M.; Xue, Z. Spatiotemporal Pattern of PM<sub>2.5</sub> Concentrations in Mainland China and Analysis of Its Influencing Factors using Geographically Weighted Regression. *Sci. Rep.* **2017**, *7*, 40607–40621. [[CrossRef](#)] [[PubMed](#)]
62. Xu, Y.; Xue, W.; Lei, Y.; Huang, Q.; Zhao, Y.; Cheng, S.; Ren, Z.; Wang, J. Spatiotemporal variation in the impact of meteorological conditions on PM<sub>2.5</sub> pollution in China from 2000 to 2017. *Atmos. Environ.* **2020**, *223*, 117215. [[CrossRef](#)]
63. Liu, Q.; Wang, S.; Zhang, W.; Li, J.; Dong, G. The effect of natural and anthropogenic factors on PM<sub>2.5</sub>: Empirical evidence from Chinese cities with different income levels. *Sci. Total Environ.* **2019**, *653*, 157–167. [[CrossRef](#)] [[PubMed](#)]
64. Zhang, Q.; Ma, Q.; Zhao, B.; Liu, X.; Wang, Y.; Jia, B.; Zhang, X. Winter haze over North China Plain from 2009 to 2016: Influence of emission and meteorology. *Environ. Pollut.* **2018**, *242*, 1308–1318. [[CrossRef](#)] [[PubMed](#)]
65. Xu, G.; Ren, X.; Xiong, K.; Li, L.; Bi, X.; Wu, Q. Analysis of the driving factors of PM<sub>2.5</sub> concentration in the air: A case study of the Yangtze River Delta, China. *Ecol. Indic.* **2020**, *110*, 105889. [[CrossRef](#)]
66. Lin, G.; Fu, J.; Jiang, D.; Wang, J.; Wang, Q.; Dong, D. Spatial variation of the relationship between PM<sub>2.5</sub> concentrations and meteorological parameters in China. *BioMed Res. Int.* **2015**, *2015*, 1–15.
67. Li, X.; Ma, Y.; Wang, Y.; Liu, N.; Hong, Y. Temporal and spatial analyses of particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>) and its relationship with meteorological parameters over an urban city in northeast China. *Atmos. Res.* **2017**, *198*, 185–193. [[CrossRef](#)]
68. Hien, P.D.; Bac, V.T.; Tham, H.C.; Nhan, D.D.; Vinh, L.D. Influence of meteorological conditions on PM<sub>2.5</sub> and PM<sub>2.5–10</sub> concentrations during the monsoon season in Hanoi, Vietnam. *Atmos. Environ.* **2002**, *36*, 3473–3484. [[CrossRef](#)]
69. Trinh, T.T.; Trinh, T.T.; Le, T.T.; Tu, B.M. Temperature inversion and air pollution relationship, and its effects on human health in Hanoi City, Vietnam. *Environ. Geochem. Health* **2019**, *41*, 929–937. [[CrossRef](#)]
70. Mažeikis, A. Urbanization influence on meteorological parameters of air pollution: Vilnius case study. *Baltica* **2013**, *26*, 51–56. [[CrossRef](#)]
71. Megaritis, A.G.; Fountoukis, C.; Charalampidis, P.E.; Denier Van Der Gon, H.A.C.; Pilinis, C.; Pandis, S.N. Linking climate and air quality over Europe: Effects of meteorology on PM<sub>2.5</sub> concentrations. *Atmos. Chem. Phys.* **2014**, *14*, 10283–10298. [[CrossRef](#)]
72. Zhou, W.; Tie, X.; Zhou, G.; Liang, P. Possible effects of climate change of wind on aerosol variation during winter in Shanghai, China. *Particuology* **2015**, *20*, 80–88. [[CrossRef](#)]
73. Cheng, X.; Zhao, T.; Gong, S.; Xu, X.; Han, Y.; Yin, Y.; Tang, L.; He, H.; He, J. Implications of East Asian summer and winter monsoons for interannual aerosol variations over central-eastern China. *Atmos. Environ.* **2016**, *129*, 218–228. [[CrossRef](#)]
74. Wang, C.; An, X.; Zhang, P.; Sun, Z.; Cui, M.; Ma, L. Comparing the impact of strong and weak East Asian winter monsoon on PM<sub>2.5</sub> concentration in Beijing. *Atmos. Res.* **2019**, *215*, 165–177. [[CrossRef](#)]
75. Backes, A.M.; Aulinger, A.; Bieser, J.; Matthias, V.; Quante, M. Ammonia emissions in Europe, part II: How ammonia emission abatement strategies affect secondary aerosols. *Atmos. Environ.* **2016**, *126*, 153–161. [[CrossRef](#)]
76. Yu, L.; Wang, G.; Zhang, R.; Zhang, L.; Song, Y.; Wu, B.; Li, X.; An, K.; Chu, J. Characterization and source apportionment of PM<sub>2.5</sub> in an urban environment in Beijing. *Aerosol Air Qual. Res.* **2013**, *13*, 574–583. [[CrossRef](#)]

77. Wu, J.; Xie, W.; Li, W.; Li, J. Effects of urban landscape pattern on PM<sub>2.5</sub> Pollution-A Beijing Case Study. *PLoS ONE* **2015**, *10*, e0142449. [[CrossRef](#)]
78. Vayenas, D.V.; Takahama, S.; Davidson, C.I.; Pandis, S.N. Simulation of the thermodynamics and removal processes in the sulfate-ammonia-nitric acid system during winter: Implications for PM<sub>2.5</sub> control strategies. *J. Geophys. Res. Atmos.* **2005**, *110*, 1–11. [[CrossRef](#)]
79. Ye, Z.; Guo, X.; Cheng, L.; Cheng, S.; Chen, D.; Wang, W.; Liu, B. Reducing PM<sub>2.5</sub> and secondary inorganic aerosols by agricultural ammonia emission mitigation within the Beijing-Tianjin-Hebei region, China. *Atmos. Environ.* **2019**, *219*, 116989. [[CrossRef](#)]
80. Gu, B.; Sutton, M.; Chang, S.; Ge, Y.; Chang, J. Agricultural ammonia emissions contribute to China's urban air pollution. *Front. Ecol. Environ.* **2014**, *12*, 265–266. [[CrossRef](#)]
81. Wu, Y.; Gu, B.; Erisman, J.W.; Reis, S.; Fang, Y.; Lu, X.; Zhang, X. PM<sub>2.5</sub> pollution is substantially affected by ammonia emissions in China. *Environ. Pollut.* **2016**, *218*, 86–94. [[CrossRef](#)] [[PubMed](#)]
82. Tao, J.; Gao, J.; Zhang, L.; Zhang, R.; Che, H.; Zhang, Z.; Lin, Z.; Jing, J.; Cao, J.; Hsu, S.C. PM<sub>2.5</sub> pollution in a megacity of Southwest China: Source apportionment and implication. *Atmos. Chem. Phys.* **2014**, *14*, 8679–8699. [[CrossRef](#)]
83. Wu, S.-P.; Zhang, Y.-J.; Schwab, J.J.; Li, Y.-F.; Liu, Y.-L.; Yuan, C.-S. High-resolution ammonia emissions inventories in Fujian, China, 2009–2015. *Atmos. Environ.* **2017**, *162*, 100–114. [[CrossRef](#)]
84. Huang, R.J.; Zhang, Y.; Bozzetti, C.; Ho, K.F.; Cao, J.J.; Han, Y.; Daellenbach, K.R.; Slowik, J.G.; Platt, S.M.; Canonaco, F.; et al. High secondary aerosol contribution to particulate pollution during haze events in China. *Nature* **2014**, *514*, 218–222. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).